

PREPARED FOR SUBMISSION TO JHEP

MAPHY-AvH/2013-01

Resummation of infrared logarithms in de Sitter space via Dyson–Schwinger equations: the ladder-rainbow approximation

A. Youssef,^a D. Kreimer^a

^a*Institut für Mathematik und Institut für Physik, Humboldt-Universität zu Berlin
Johann von Neumann-Haus, Rudower Chaussee 25, 12489 Berlin, Germany*

E-mail: youssef@mathematik.hu-berlin.de,
kreimer@mathematik.hu-berlin.de

ABSTRACT: We study the infrared (large separations) behavior of a massless minimally coupled scalar QFT with a quartic self-interaction in de Sitter. We show that the perturbation series in the interaction strength λ is secular, i.e it does not lead to a uniform approximation of the solution in the infrared region. Only a non-perturbative resummation can capture the correct IR behavior. We seek to justify this picture using the Dyson–Schwinger equations in the ladder-rainbow approximation. We are able to write down an ordinary differential equation obeyed by the two-point function and perform its asymptotic analysis. Opposing a widespread belief that IR divergences are regulated through a dynamical mass generation, our work suggest an even stronger IR decay than the power-law one resulting from a mass term.

Contents

1	Introduction	2
1.1	Prelude	2
1.2	De Sitter geometry and QFT	3
1.3	The massless field	3
2	Dyson–Schwinger equations	5
3	Solution of the master equation	8
4	Dynamical mass generation?	9
5	Discussion	9
A	Conventions	11
B	Flat space master equation	11
C	Local analysis	11
C.1	Frobenius series at $z = 0$	12
C.2	Asymptotic behavior near $z = 1$	12
C.3	Asymptotic behavior near $z \rightarrow \infty$	13
D	WKB approximation	14

Contents

1 Introduction

1.1 Prelude

The two-point function Δ_{m^2} of a massive scalar field in de Sitter decays in the IR region, i.e for largely separated spacetime points. For small masses, Δ_{m^2} develops a $1/m^2$ pole, making the massless limit ill-defined and the quantization of the theory a non-trivial task. This situation is to contrast with the flat space case, where massive and massless fields two-point functions decay for large invariant distance μ (as $\mu^{-2}e^{-m^2\mu}$ and μ^{-2} respectively).

Several propositions to quantize the massless minimally coupled (mmc) field co-exist in the literature, depending on how one interprets and cures this ill-defined massless limit. The first and most popular direction is to understand the problematic massless limit as an indication of the impossibility of maintaining the full de Sitter invariance of the quantum theory. Indeed, breaking de Sitter invariance leads to a well-defined quantization and to interesting cosmological implications [1, 2].

Another option, investigated in [3, 4] (see also the earlier Starobinsky’s stochastic approach [5, 6]), is to recognize the necessity of the inclusion of a small (say quartic) interaction of strength λ . The idea being that the presence of a small but non-vanishing λ will make the massless limit smooth (in other terms, the limits $m \rightarrow 0$ and $\lambda \rightarrow 0$ do not commute). These investigations lead to the conclusion that interactions induce a dynamically generated mass $m_{\text{eff}}^2 \propto \sqrt{\lambda}H$ -where H is the Hubble constant-that regulates the IR behavior. In particular, in [4] an interesting exact treatment of the zero-mode in the interacting theory (which amounts to performing a non-gaussian but one dimensional integral) is given, and the author argues in favor of a constant massless two-point function proportional to $1/\sqrt{\lambda}$. This approach, as recently shown in [7] leads again to the conclusion that interactions induce a dynamically generated mass.

Yet another approach can be found in [8, 9], where a “renormalized” two-point function Δ_0 is defined through the subtraction of the divergent $1/m^2$ term. This approach leads to a well-defined two-point function Δ_0 , which is growing in the IR. The construction of Δ_0 presents a lot of advantages, as it preserves causality and positivity. We have also shown in [10] that it leads to the observed scale-invariant CMB power-spectrum. Our present work is based on this free propagator.

While we indeed believe that the inclusion of an interaction, whatever small, has drastic consequences on the quantization of the massless field and its IR behavior, we argue here in favor of a quite different overall picture. The free propagator Δ_0 , as well as its perturbative corrections are growing. However, when the leading IR contributions are summed up, the full non-perturbative propagator G is decaying. In other words, the perturbative expansion develop secular terms, making the perturbative expansion non-uniform in the IR. Contrary to the previously discussed papers, we suggest that the decay of G is too rapid to be interpreted as resulting from a dynamical mass generation. We take a first step to justify this picture by summing, via a linear Dyson–Schwinger equation, the so called ladder-rainbow diagrams.

The organization of the paper is as follows. After a rapid introduction to the basics of de Sitter geometry and QFT, we write down in section II the Dyson–Schwinger equations in the so-called ladder-rainbow approximation. Taking full-advantage of de Sitter symmetry,

we are able to reduce this integral equation into an ordinary differential equation. The asymptotic analysis of which is studied in section III. A concise discussion of dynamical mass generation is given in section IV. Finally, section V is a discussion of several important points.

1.2 De Sitter geometry and QFT

The d -dimensional de Sitter spacetime can be identified with the real one-sheeted hyperboloid in the $d + 1$ Minkowski spacetime M_{d+1} :

$$X_d = \left\{ x \in \mathbb{R}^{d+1}, x^2 = -R^2 \right\}$$

with $R = H^{-1} > 0$ being the de Sitter “radius”. This definition of the de Sitter manifold reveals the maximal symmetry of X_d under the action of the de Sitter group $SO_0(1, d)$. Let $\mu(x, x')$ denote the geodesic distance between the points x and x' . It is useful to introduce the quantity z given for spacelike separations ($\mu^2 > 0$ or $0 < z < 1$) by $z = \cos^2\left(\frac{\mu}{2R}\right)$. As a general rule, we will work in the spacelike region and then analytically continue the propagators. As shown in [11], when acting on functions of the invariant $\mu(x, x')$, the Laplace-Beltrami operator reduces to the ordinary differential operator :

$$-\square = \frac{d^2}{d\mu^2} + (d-1)A(\mu)\frac{d}{d\mu} \quad (1.1)$$

where

$$\begin{cases} A(\mu) = \mu^{-1}, & \text{on Minkowski space} \\ A(\mu) = \frac{\cot(\mu/R)}{R}, & \text{on de Sitter space.} \end{cases}$$

In terms of the variable z , this becomes the hypergeometric operator

$$-\square = \mathbf{H} = z(1-z)\frac{d^2}{dz^2} + [c - (a+b+1)z]\frac{d}{dz} - ab$$

where

$$a = \frac{d-1 + \sqrt{(d-1)^2 - 4m^2R^2}}{2}, \quad b = \frac{d-1 - \sqrt{(d-1)^2 - 4m^2R^2}}{2}, \quad c = d/2.$$

In particular, in the massless case we have

$$\mathbf{H} = z(1-z)\frac{d^2}{dz^2} + \frac{d}{2}(1-2z)\frac{d}{dz}. \quad (1.2)$$

1.3 The massless field

The physical reason behind the appearance of strong IR effects in de Sitter can be simply understood: the rapid expansion of the spacetime dilate correlation patterns. After all this is the exact reason why a de Sitter inflationary phase in the early universe solves many problems of the hot big-bang model. These IR effects are naturally stronger for massless (and non-conformally invariant) fields such as the massless minimally coupled (mmc)¹ and the graviton. We review here these IR divergences in the mmc case.

¹This means that there is no term in the action proportional to $\mathcal{R}\phi^2$, $\mathcal{R}$ being the Ricci scalar, which is constant in de Sitter spacetime

Recall that in the Bunch-Davies vacuum state the massive propagator for a scalar field reads :

$$\Delta(x, x') = \Gamma(-\sigma) \frac{\Gamma(\sigma + d - 1)}{(4\pi)^{d/2} R^{d-2} \Gamma(\frac{d}{2})} {}_2F_1 \left[-\sigma, \sigma + d - 1, \frac{d}{2}; z \right] \quad (1.3)$$

where $\sigma = -\frac{d}{2} + \sqrt{\frac{d^2}{4} - m^2 R^2}$ and ${}_2F_1$ is the hypergeometric function. In the massless limit, $\sigma \rightarrow 0$, this expression diverges because of the pole of the first Gamma function and we have the small mass expansion:

$$\Delta(x, x') \sim \frac{d}{4\pi^{\frac{d+1}{2}}} \Gamma\left(\frac{d-1}{2}\right) \frac{1}{m^2 R^d} + \text{regular terms in } m.$$

Note that in the flat space limit ($R \rightarrow \infty$), this singular term is absent and the massless limit is smooth². More precisely we have (for $d > 2$):

$$\Delta^{\text{flat}}(x, x') = -\frac{i\pi}{4\pi^{d/2}} \left(\frac{m^2}{\mu^2}\right)^{\frac{d-2}{4}} H_{\frac{d}{2}-1}^{(2)}\left(2\sqrt{m^2\mu^2}\right) \underset{m \rightarrow 0}{\sim} \left(\frac{1}{2\sqrt{\pi}}\right)^d \Gamma\left(\frac{d}{2} - 1\right) \mu^{2-d}$$

where $\mu = \sqrt{(x - x')^2}$ is the invariant distance and $H^{(2)}$ is the Hankel function of the second kind.

One of the first papers studying the mmc scalar field in de Sitter is [1], where the authors prove that a usual de Sitter-invariant Fock space quantization is impossible in this case. They then propose to trade the de Sitter $\text{SO}(1, d)$ invariance for a smaller one, say a $\text{SO}(d)$ invariance. Equivalently, it is a common belief among workers in the field that the scale-invariant power-spectrum leads necessarily to a breakdown of de Sitter invariance and that some physical quantities might thus become time-dependent. Several authors later proposed different treatments of the mmc field, among which [9] is one of the most exhaustive. Here the divergent term is subtracted, a “renormalized” de Sitter-invariant two-point function is computed³ and it reads (we work in $d = 4$ for simplicity):

$$\Delta_0(z) = \frac{1}{(4\pi)^2 R^2} \left[\frac{1}{1-z} - 2 \ln(1-z) \right]. \quad (1.4)$$

In [10] we also proved that Δ_0 gives the observed scale-invariant power spectrum. We will use this propagator for the mmc hereafter (and we will omit the subscript 0 when no confusion can arise).

²This means that the flat space limit ($R \rightarrow \infty$) and the massless limit ($m \rightarrow 0$) do not commute. This is a physically important fact and might mean that even a small amount of curvature - like in today’s universe - might have important consequences on massless fields.

³The draw-back is that the two-point function no longer verifies the equation of motion $\square\phi = 0$, instead it verifies the anomalous equation:

$$\square\phi = -\frac{3}{8\pi^2}.$$

This simple renormalization procedure has been used implicitly in several earlier works. However, the major contribution of [9] is proving that on a suitably chosen subspace of states $\mathcal{E}$, the equation of motion is effectively restored. This “Physical” space of states should be regarded the same way as we regard the one that appears in the quantization of gauge theory (for instance the space of transverse photons in QED). Moreover, the authors were able to show that the renormalized two-point function defines a positive kernel when restricted to $\mathcal{E}$, thus enabling a probabilistic interpretation of the theory.

2 Dyson–Schwinger equations

The Dyson–Schwinger equations, an infinite set of integral equations between the n -point functions, are the equations of motion of QFT. These equations, through some truncation schemes, give a convenient handle on non-perturbative effects in the theory. From now on we study the QFT given by the action:

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} g_{\mu\nu} \partial^\mu \phi \partial^\nu \phi - \frac{1}{2} m^2 \phi^2 - \frac{\lambda}{4!} \phi^4 \right],$$

where $dV_x = \sqrt{-g} d^4x$ is the invariant volume element.

The (unrenormalized) Dyson–Schwinger equations for the exact propagator involve the four-point function Γ_4 :

$$G(x, x') = \Delta(x, x') + \int dV_{y_1} dV_{y_2} \Delta(x, y_1) \Sigma(y_1, y_2) G(y_2, x') \quad (2.1)$$

$$\Sigma(x, y) = \frac{1}{2} (-i\lambda)^2 \int dV_{z_1} dV_{z_2} dV_{z_3} G(x, z_1) G(x, z_2) G(x, z_3) \Gamma_4(z_1, z_2, z_3, y) \quad (2.2)$$

where G is the exact Feynman propagator, Δ the free one, Σ the self-energy and Γ_4 the exact 4-point function. A graphical representation of these two-equations is given in figure (1). The first equation is known as the Dyson equation which in flat spacetime is readily solved in Fourier space and is equivalent to the usual geometric series of self-energies:

$$G(p) = \frac{\Delta(p)}{1 - \Delta(p)\Sigma(p)}.$$




Figure 1. Dyson–Schwinger equation relating the two and four point functions.

Actually, let us now analyze the Dyson–Schwinger equations for our purposes in some detail. We concentrate on the flat space case, the results being immediately generalizable to de Sitter spacetime. To start with, we note that our self-energy starts at two loops. This is obvious, as also in de Sitter the two-point tadpole graph has zero renormalized value, the tadpole vanishes [12]. Thus, the expansion of the self-energy starts at two loops, with the first term in the expansion indeed furnishing a primitive element in the Hopf algebra H of ϕ^4 theory, which provides a Hochschild one-cocycle B_+ . Omitting higher Hochschild co-cycles -which seems reasonable as they do not alter the algebraic structure of the Dyson–Schwinger equations-, we find the combinatorial Dyson–Schwinger equation for the self-energy

$$X(\lambda^2) = 1 - \frac{\lambda^2}{6} B_+ \left(\frac{1}{X^3(\lambda^2)} \right).$$




Figure 2. There is no contribution from this graph.

Its fix-point is a formal series $\in H[[\lambda^2]]$ such that the application of renormalized Feynman rules Φ_R gives $\Phi_R(X(\lambda^2)) = 1 - \Sigma$.

Hochschild cohomology ensures that we renormalized by local counterterms, and ensures that each contributing graph is divided by its symmetry factor, as it should [13, 14].

We drastically simplify this system by linearizing it to a commutative and co-commutative Hopf algebra generated from simple concatenations of the cocycle B_+ .

$$X(\lambda^2) = 1 - \frac{\lambda^2}{6} B_+(X(\lambda^2)).$$

Graphically, the two equations read as in Fig. 3.




Figure 3. Flat space non-linear and linear Dyson–Schwinger equations.

Explicitly, the two expansions in graphs read as in Fig.4. Note that in the lin-

$$\begin{aligned}
\text{---}\bullet\text{---} &= \text{---} - \left(\frac{\lambda^2}{6} \text{---}\bigcirc\text{---} + \frac{\lambda^4}{12} \text{---}\bigcirc\!\!\bigcirc\text{---} + \frac{\lambda^6}{24} \text{---}\bigcirc\!\!\bigcirc\!\!\bigcirc\text{---} + \frac{\lambda^6}{72} \text{---}\bigcirc\!\!\bigcirc\!\!\bigcirc\!\!\bigcirc\text{---} + \mathcal{O}(\lambda^8)\right) \\
\text{---}\bullet\text{---} &= \text{---} - \left(\frac{\lambda^2}{6} \text{---}\bigcirc\text{---} + \frac{\lambda^4}{36} \text{---}\bigcirc\!\!\bigcirc\text{---} + \frac{\lambda^6}{216} \text{---}\bigcirc\!\!\bigcirc\!\!\bigcirc\text{---} + \mathcal{O}(\lambda^8)\right)
\end{aligned}$$

Figure 4. Non-linear and linear Dyson–Schwinger equations expanded out in perturbation theory. Note that the linear Dyson–Schwinger equations misses some graphs, which also results in wrong symmetry factors. It nevertheless gives a strict subset of the full expansion at any order.

earized case, graphs do not contribute by their symmetry factors any more, as “insertion places” are missing. More precisely, in the so-called ladder-rainbow approximation we have $\Gamma_4(z_1, z_2, z_3, y) = \delta(y, z_1)\delta(y, z_2)\delta(y, z_3)$ where the distribution $\delta(x, y) = \frac{1}{\sqrt{-g}}\delta(x - y)$ is defined via

$$\int dV_y \delta(x, y) f(y) = f(x).$$

Moreover, in this approximation the self-energy is given by

$$\Sigma(x, y) = \frac{1}{6}(-i\lambda)^2 G(x, y) \Delta^2(x, y).$$

As stated above, we also consider only the linearized version of the Dyson equation by replacing the exact propagator by the free one in the right hand side of (2.1). This means in particular that we do not sum over all self-energy chains. Finally we obtain the closed linear integral equation for the two-point function

$$G(x, x') = \Delta(x, x') - \frac{\lambda^2}{6} \int dV_{y_1} dV_{y_2} \Delta(x, y_1) G(y_1, y_2) \Delta^2(y_1, y_2) \Delta(y_2, x') \quad (2.3)$$

which can be transformed into the differential equation [20]:

$$(\square_x + m^2)(\square_{x'} + m^2)G(x, x') - \frac{\lambda^2}{6}\Delta^2(x, x')G(x, x') = -i(\square_x + m^2)\delta(x, x').$$

For maximally symmetric spacetimes and vacuum states this reduces to:

$$\left[(\square + m^2)^2 - \frac{\lambda^2}{6}\Delta^2(x, x') \right] G(x, x') = -i(\square + m^2)\delta(x, x'). \quad (2.4)$$

In terms of the z variable and the hypergeometric operator $\mathbf{H}$, using the formulas (1.1), the homogeneous ⁴ part of the preceding equation becomes:

$$\boxed{[\mathbf{H}^2 - \epsilon \Delta^2] G(z) = 0, \quad \epsilon = \lambda^2/6.} \quad (2.5)$$

Note that for the non-linear Dyson–Schwinger equation, the corresponding differential equation would become highly non-linear even in flat space -see [15]-, and on top of that

⁴The inhomogeneity proportional to a delta function will be responsible to a shift that can be absorbed in the integration constants

non-linearity we would have all the complexities of the hypergeometric operator on which we now focus. All further discussions of the non-linear case are postponed to future work.

This ordinary differential equation is the central equation of our work and we will refer to it as the master equation in what follows. We will use the following boundary conditions, which are the natural generalization of the Bunch-Davies vacuum state (a direct derivation of these conditions from the integral equation will be given elsewhere): regularity for antipodal points (i.e at $z = 0$) and a flat spacetime singularity at short distances (at $z \rightarrow 1$). The flat space singularity is proportional to (see Appendix B):

$$\mu^{-\frac{1}{2}} \sqrt{\frac{\sqrt{\epsilon+64\pi^2}}{\pi^2} + 8}. \quad (2.6)$$

Developing the relation $z = \cos^2\left(\frac{\mu}{2R}\right)$ near $\mu \rightarrow 0$, we have the asymptotic behavior of G at $z = 1$:

$$G \underset{z \rightarrow 1}{\sim} a (1-z)^\nu, \quad \nu = -\frac{1}{4} \sqrt{\frac{\sqrt{\epsilon+64\pi^4}}{\pi^2} + 8}.$$

where a is a given constant independent of ϵ .

3 Solution of the master equation

We study the homogeneous master equation (2.5), which we reproduce here explicitly for convenience:

$$\begin{aligned} G^{(4)} + \left(\frac{6}{z} + \frac{6}{z-1}\right) G^{(3)} + \left(\frac{6}{z^2} - \frac{22}{z} + \frac{22}{z-1} + \frac{6}{(z-1)^2}\right) G'' \\ + \left(\frac{8}{(z-1)^2} - \frac{8}{z^2}\right) G' - \frac{\epsilon}{256\pi^4(1-z)^2 z^2} \left(\frac{1}{1-z} - 2\ln(1-z)\right)^2 G = 0. \end{aligned} \quad (3.1)$$

This equation possesses one regular singular point at $z = 0$ and two irregular singular points at $z = 1$ and $z = \infty$. The local study performed around these points in Appendix C shows that the interaction term (proportional to ϵ) is negligible near $z = 0$ and is very important near $z = \infty$.

We then use a WKB approximation (or Liouville-Green for mathematicians) to obtain global approximations to the four fundamental solutions of (C.1). The solution that posses the correct $z \rightarrow 1$ has the following IR behavior:

$$G \sim \frac{1}{z^{3/2} \ln^{\frac{3}{4}}(-z)} \exp \left[- \left(\frac{\sqrt[4]{\epsilon} \ln^{\frac{3}{2}}(-z)}{3\sqrt{2}\pi} + \frac{9\pi\sqrt{\ln(-z)}}{\sqrt{2}\sqrt[4]{\epsilon}} \right) \right].$$

As the other solutions have a sub-leading contribution near $z = 1$, they (notably the one with $\omega = 1$ in the appendix C) can spoil the IR behavior by making it grow instead of decaying. This will be checked via a rigorous analysis elsewhere.

We note that, while the approximation near $z = 1$ obtained via a certain WKB approximation is only valid for strong couplings $\epsilon \ll 1$, the asymptotic behavior obtained via local analysis (see Appendix C) is exact for any ϵ .

Finally an important remark: the phenomenon described here belongs to a large class of phenomena in mathematics and physics known as secular perturbation theory. The simplest such example is given by the boundary value problem [16]:

$$y'' + \epsilon y' + y = 0, \quad \text{for } t > 0, \quad y(0) = 0, y'(0) = 1.$$

To the first order in perturbation theory in ϵ the solution is unbounded for large t :

$$y(t) = \sin(t) - \frac{1}{2}\epsilon t \sin(t).$$

Unlike this perturbative solution, the exact solution is bounded:

$$y(t) = \frac{1}{\sqrt{1-\epsilon^2/4}} e^{-\epsilon t/2} \sin\left(t \sqrt{1-\epsilon^2/4}\right).$$

In the case under consideration, the multiple scales method [16, 17] (consisting of introducing a new scale $\tau = \epsilon t$, treat it as independent of t , and use this freedom to kill the secular terms) yields, to the first iteration, the uniform approximation

$$y(t) \sim e^{-\epsilon t/2} \sin(t).$$

We will investigate the applicability of these multiple scales methods to the study of strong IR effects in de Sitter in a future publication. Such secular behavior is usually related to a resonance in the system. Accordingly, we suggest that the expansion of the de Sitter background plays the role of a resonant external coupling.

4 Dynamical mass generation?

The authors of [7] retrieve the dynamically generated mass predicted by Starobinsky in [5]. Their approach is based on the constant “free” two-point function obtained in [4]. We show in this section how this result can be readily retrieved through our master equation.

If we consider the constant two-point function $\Delta_0 \propto \frac{1}{\sqrt{\lambda}}$ given in [4] (note that it is not really a free two-point function as it diverges for $\lambda \rightarrow 0$), our master equation (2.5) summing the ladder-rainbow diagrams has the four IR behaviors at $z \rightarrow \infty$:

$$\left\{ z^{\frac{1}{2}(-\sqrt{9-4\lambda\Delta_0}-3)}, z^{\frac{1}{2}(\sqrt{9-4\lambda\Delta_0}-3)}, z^{\frac{1}{2}(-\sqrt{9+4\lambda\Delta_0}-3)}, z^{\frac{1}{2}(\sqrt{9+4\lambda\Delta_0}-3)} \right\}.$$

The boundary conditions select $z^{\frac{1}{2}(\sqrt{9-4\lambda\Delta_0}-3)}$. As a mass m field two-point function (1.3) decays like $z^{\frac{1}{2}(\sqrt{9-4m^2}-3)}$, this implies a dynamically generated mass $m_d^2 = \lambda\Delta_0 \propto \sqrt{\lambda}$.

5 Discussion

We end our work with a concise discussion of several important issues:

- A crucial technical point in our analysis is that we are able to take full-advantage of de Sitter-symmetry by transforming the Dyson–Schwinger integral equation into an ordinary differential equation (depending only on the scalar variable z).
- It is important to stress that our conclusions were drawn in a very restrictive context, namely the ladder-rainbow approximation, which is a drastic simplification of the full theory. A better control of this approximation has to be achieved, mainly by summing over a larger family of diagrams (for instance using the non-linear Dyson–Schwinger equation discussed before). We note however that the difficulty to rigorously justify the used approximation is a general trend in hard problems such as non-perturbative QFT, a classical example being the study of bound-states using the Bethe-Salpeter equation [21].

- We have been able to avoid any discussion about UV-renormalization because we transform the integration over loops into differential equations. Hence renormalization only intervenes through the integration constants. This is actually the generic situation for the study of solutions of Dyson–Schwinger equations: for a kinetic renormalization scheme, renormalization condition amount to fixing the boundary conditions of the equations.
- The secularity of perturbation theory exists already in flat space. However, unlike the de Sitter case, it does not change the IR behavior of the two-point function, because of a rapidly decaying overall μ^{-2} factor. Indeed the ladder-rainbow two-point function in flat space reads (see appendix B):

$$\mu^{-\sqrt{2+\frac{\sqrt{\epsilon+64\pi^4}}{4\pi^2}}} = \frac{1}{\mu^2} \left[1 - \frac{\epsilon \ln \mu}{256\pi^4} + \frac{\epsilon^2 (5 \ln \mu + 2 \ln^2 \mu)}{262144\pi^8} + \dots \right].$$

- The Fourier transform and power-spectra of the non-perturbative functions G has to be studied.
- The IR behavior of the graviton field in de Sitter spacetime is one of the most important open questions in cosmology. We believe that the non-perturbative effects exhibited in the present work for the mmc scalar field can illuminate this issue. On top of these non-perturbative effects, we also expect an interesting interplay with non-trivial gauge artifacts similar to the ones already exhibited for the photon field in de Sitter in [18].
- In our study we neglect the back-reaction of the quantum fields on the background metric. However the fact that the interacting two-point function G decays in the IR is of paramount relevance to the difficult but crucial issue of whether the back-reaction can be neglected. Ultimately, this is equivalent to understanding the stability of the quantum de Sitter spacetime.

Acknowledgments

D. Kreimer is supported by the Alexander von Humboldt Foundation and the BMBF through an Alexander von Humboldt Professorship.

A Conventions

We use the conventions of [19]. The metric is the mostly minus one. Consider the action

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} g_{\mu\nu} \partial^\mu \phi \partial^\nu \phi - \frac{1}{2} m^2 \phi^2 - \frac{\lambda}{4!} \phi^4 \right]$$

where g is the determinant of the metric. The free Feynman propagator $\langle T\phi(x)\phi(x') \rangle = \Delta(x, x')$ verifies:

$$(\square_x + m^2)\Delta(x, x') = -i \frac{\delta(x - x')}{\sqrt{-g}}.$$

The conventions for the Fourier transform are

$$\Delta(x) = \int \frac{d^4k}{(2\pi)^4} e^{-ikx} \Delta(k)$$

Feynman rules are given by:

Propagator:	$\Delta(x, x')$
Vertex:	$-i\lambda_B$
Mass renormalization:	$-i(m_B - m^2)$
Field renormalization:	$-i(Z - 1)\square_x$

B Flat space master equation

In the flat space the master equation (2.5), with $A = 1/\mu$ and $\Delta(\mu) = \frac{1}{4\pi^2\mu^2}$ is readily solved and gives the four fundamental solutions with

$$\mu^\rho, \quad \rho = \pm \sqrt{2 \pm \frac{\sqrt{\epsilon + 64\pi^4}}{4\pi^2}}.$$

The boundary conditions, included in the integral equation for instance, select the solution for which $\rho = -\sqrt{2 + \frac{\sqrt{\epsilon + 64\pi^4}}{4\pi^2}}$.

C Local analysis

We study the homogeneous master equation (2.5), which we reproduce here explicitly for convenience:

$$\begin{aligned} G^{(4)} + \left(\frac{6}{z} + \frac{6}{z-1} \right) G^{(3)} + \left(\frac{6}{z^2} - \frac{22}{z} + \frac{22}{z-1} + \frac{6}{(z-1)^2} \right) G'' \\ + \left(\frac{8}{(z-1)^2} - \frac{8}{z^2} \right) G' - \frac{\epsilon}{256\pi^4(1-z)^2 z^2} \left(\frac{1}{1-z} - 2\ln(1-z) \right)^2 G = 0. \end{aligned} \quad (\text{C.1})$$

This equation possesses one regular singular point at $z = 0$ and two irregular singular points at $z = 1$ and $z = \infty$. We perform here a local analysis near these points.

C.1 Frobenius series at $z = 0$

The indicial polynomial is $P_0(\lambda) = \lambda^2(\lambda^2 - 1)$. Our equation being fourth-order and the roots of P_0 being separated by integers, some care has to be taken in the application of the Frobenius method (see [22] for the explicit algorithm). We obtain the local behaviors:

$$\left\{ \frac{1}{z}, \ln z, 1, z \right\}$$

More precisely the first terms of the four solutions are given by:

$$\begin{cases} G_1(z) \sim z + \frac{2z^2}{3} + \left(\frac{\epsilon}{18432\pi^4} + \frac{5}{9} \right) z^3 \sim z, \\ G_2(z) \sim \frac{1}{z} + 84 - \left(216 + \frac{15\epsilon}{512\pi^4} \right) z + \left[-48 + \left(36 + \frac{3\epsilon}{256\pi^4} \right) z \right] \ln z \sim \frac{1}{z}, \\ G_3(z) \sim 1 - z + \left(-\frac{2}{3} + \frac{\epsilon}{3072\pi^4} \right) z^2 \sim 1, \\ G_4(z) \sim 9z + \left(\frac{40}{9} - \frac{7\epsilon}{4608\pi^4} \right) z^2 + \left[2 - 2z + \left(-\frac{4}{3} + \frac{\epsilon}{1536\pi^4} \right) z^2 \right] \ln z \sim 2 \ln z. \end{cases}$$

This analysis implies that, the boundary condition imposing regularity at $z = 0$ eliminates 2 of the 4 solutions.

C.2 Asymptotic behavior near $z = 1$

$\epsilon = 0$. In this case the point $z = 1$ is a regular singular point. The free master equation being invariant under the transformation $z \rightarrow 1 - z$, the asymptotic behaviors at $z = 1$ can be directly obtained from the behavior near $z = 0$ (also the equation is then exactly solvable) and are given by:

$$\left\{ 1 - z, \frac{1}{1 - z}, \ln(1 - z), 1 \right\}.$$

$\epsilon > 0$. Even if the point $z = 1$ is an irregular singular point, this “irregularity” appears only at the first order when $z \rightarrow 1$. At zeroth-order in $1 - z$, one can thus still define an indicial polynomial which is given by:

$$P_1(\lambda) = \lambda^4 - \lambda^2 - \frac{\epsilon}{256\pi^4}.$$

The roots of which read:

$$\lambda = \pm \frac{1}{4} \sqrt{\frac{\sqrt{\epsilon + 64\pi^4}}{\pi^2}} \pm 8$$

and are not separated by integers for generic ϵ . The leading behaviors near $z = 1$ is thus given by

$$(1 - z)^\lambda, \quad \lambda = \pm \frac{1}{4} \sqrt{\frac{\sqrt{\epsilon + 64\pi^4}}{\pi^2}} \pm 8$$

We can verify that this is the full leading behavior at $z = 1$.

C.3 Asymptotic behavior near $z \rightarrow \infty$

By making the ansatz $G = e^S$ and using the dominant balance method recursively [17], we are able to derive the leading behavior near infinity:

$$G_\omega \sim \frac{1}{z^{3/2} \ln^{3/4}(-z)} \exp \left[\left(\frac{\omega \sqrt[4]{\epsilon} \ln^{3/2}(-z)}{3\sqrt{2}\pi} + \frac{9\pi \sqrt{\ln(-z)}}{\sqrt{2}\omega \sqrt[4]{\epsilon}} \right) \right], \quad \omega^4 = 1.$$

Obtaining this result, necessitates some lengthy calculations that we choose not to reproduce here.

D WKB approximation

Consider the general fourth-order differential equation (of course the parameter δ can be introduced in many different ways, leading to more or less accurate approximations):

$$\delta^4 \left[G^{(4)} + a_3 G^{(3)} + a_2 G'' + a_1 G' \right] - a_0 G = 0.$$

When $\delta \rightarrow 0$, each of the four elementary solutions in the WKB approximation read (in our situation, $\delta = \epsilon^{-1/4}$, meaning that we perform a strong coupling expansion):

$$G \sim \exp \left[\frac{1}{\delta} (S_0 + \delta S_1 + \delta^2 S_2) \right]$$

where

$$\begin{aligned} S_0 &= \omega \int^z \sqrt[4]{a_0}, \quad \omega^4 = 1, \\ S_1 &= \frac{-1}{8} \left(3 \log a_0 + 2 \int^z a_3 \right), \\ S_2 &= \frac{1}{128\omega} \int^z \frac{1}{\sqrt[4]{a_0}} \left[-\frac{45a_0'^2}{a_0^2} + 48a_3' + \frac{40a_0''}{a_0} + 12a_3^2 - 32a_2 \right]. \end{aligned} \tag{D.1}$$

These solutions have the asymptotic behaviors

$$\left\{ \begin{aligned} G_\omega &\sim_{z \rightarrow 1} (1-z) \frac{\omega \sqrt[4]{\epsilon}}{4\pi} + \frac{\pi}{\omega \sqrt[4]{\epsilon}}, \\ G_\omega &\sim_{z \rightarrow \infty} \frac{1}{z^{3/2} \ln^{3/4}(-z)} \exp \left[\left(\frac{\omega \sqrt[4]{\epsilon} \ln^{3/2}(-z)}{3\sqrt{2}\pi} + \frac{9\pi \sqrt{\ln(-z)}}{\sqrt{2}\omega \sqrt[4]{\epsilon}} \right) \right]. \end{aligned} \right.$$

Comparing the WKB results to the exact local analysis performed before, we find that they are exactly the same as $z \rightarrow \infty$. They also coincide near $z \rightarrow 1$ in the strong coupling limit $\epsilon \gg 1$. This is a verification of our previous calculations. The “minimal” choice corresponding to the boundary value at $z = 1$ will be $\omega = -1$, implying that the real part of the resummed two-point function decay at $z \rightarrow \infty$.

However this WKB approximation breaks down near $z = 0$, preventing us from implementing the boundary conditions at $z = 0$. This breakdown can be explicitly seen because the following relations in the expansion $G \sim \exp \left[\frac{1}{\delta} (S_0 + \delta S_1 + \delta^2 S_2) \right]$ do not hold near $z = 0$:

$$S_0 \gg S_1 \gg S_2.$$

This was also expected, since the point $z = 0$ is a degenerate regular singular point, i.e having equal exponents (see [23] p.202). The global analysis of such equations, specially higher-order ones, is a non-trivial task (see [24] and references therein) that we will examine in a future work.

References

- [1] B. Allen and A. Folacci, *The massless minimally coupled scalar field in de Sitter space*, Phys. Rev. D **35**(1987)3771.
- [2] R. Woodard, *De Sitter Breaking in Field Theory*, [arXiv:gr-qc/0408002].
- [3] J. Serreau, *Effective potential for quantum scalar fields on a de Sitter geometry*, Phys. Rev. Lett. **107**(2011)191103, [arXiv:1105.4539 [hep-th]].
- [4] A. Rajaraman, *On the proper treatment of massless fields in Euclidean de Sitter space*, Phys. Rev. D **82**(2010)123522, [arXiv:1008.1271 [hep-th]].
- [5] A. Starobinsky and J. Yokoyama, *Equilibrium State of a Massless Self-Interacting Scalar Field in the De Sitter Background*, Phys. Rev. D **50**(1994)6357.
- [6] R. Woodard, *Generalizing Starobinskii's Formalism to Yukawa Theory and to Scalar QED*, J. Phys. Conf. Ser. **68**(2007)012032, [arXiv:gr-qc/0608037].
- [7] M. Beneke, P. Moch, *On "dynamical mass" generation in Euclidean de Sitter space*, [arXiv:1212.3058].
- [8] A. Folacci, *BRST quantization of the massless minimally coupled scalar field in de Sitter space (zero modes, euclideanization and quantization)*, Phys. Rev. D **46**(1992)2553, [arXiv:0911.2064 [gr-qc]].
- [9] J. Bros, H. Epstein and U. Moschella, *Scalar tachyons in the de Sitter universe*, Lett. Math. Phys. **93**(2010)203, [arXiv:1003.1396 [hep-th]].
- [10] A. Youssef, *Do scale invariant fluctuations break de Sitter invariance?*, Phys. Lett. B **718**(2013)1095, [arXiv:1203.3171 [gr-qc]].
- [11] B. Allen and T. Jacobson, *Vector two-point functions in maximally symmetric spaces*, Commun. Math. Phys. **103**(1986)669.
- [12] D. Marolf and I. A. Morrison, *Infrared stability of de Sitter space: Loop corrections to scalar propagators*, Phys. Rev. D **82**(2010)105032, [arXiv:1006.0035 [gr-qc]].
- [13] D. Kreimer, *Algebraic Structures in local QFT*, Nucl. Phys. Proc. Suppl. **205-206**(2010)122 [arXiv:1007.0341 [hep-th]].
- [14] D. Kreimer and W. D. van Suijlekom, *Recursive relations in the core Hopf algebra*, Nucl. Phys. B **820**(2009)682 [arXiv:0903.2849 [hep-th]].
- [15] D. J. Broadhurst and D. Kreimer, *Exact solutions of Dyson-Schwinger equations for iterated one loop integrals and propagator coupling duality*, Nucl. Phys. B **600**(2001)403 [hep-th/0012146].
- [16] M.H. Holmes, *Introduction to perturbation methods*, Springer-Verlag New York (1995).
- [17] C.M. Bender and S.A. Orszag, *Advanced mathematical methods for scientists and engineers*, McGraw-Hill (1978).
- [18] A. Youssef, *Infrared behavior and gauge artifacts in de Sitter spacetime: the photon field*, Phys. Rev. Lett. **107**(2011)021101, [arXiv:1011.3755 [gr-qc]].
- [19] J. Collins, *Renormalization*, Cambridge university press (1984).
- [20] R. Delbourgo, D. Elliott and D. McAnally, *Dimensional Renormalization in ϕ^3 theory: ladders and rainbows*, Phys. Rev. D **55**(1997)5230, [arXiv:hep-th/9611150].

- [21] C. Itzykson and J-B. Zuber, *Quantum Field Theory*, Dover Publications (2006).
- [22] M. A. Barkatou, T. Cluzeau, C. El Bacha, *Frobenius Method for Computing Power Series Solutions of Linear Higher-Order Differential Systems*, Proceedings of Mathematical Theory of Networks and Systems (MTNS) 2010, hal-00633268.
- [23] F.W.J. Olver, *Asymptotics and special functions*, AKP classics (1997).
- [24] T. Koike, *Asymptotics of the spectrum of Heun's equation and the exact WKB analysis*, in Toward the exact WKB analysis of differential equations, linear or non-linear, Kyoto Univ. Press, Kyoto (2000).